



L 31461-66

ACC NR: AP6023114

solutions were degassed by the freezing method. The formation of hydrazyls in electrochemical oxidation of the original compounds can be depicted by the scheme:



Thus, the authors have shown that electrochemical oxidation as well as electrochemical reduction of compounds of the diphenylpicrylhydrazine type lead to the formation of free radicals, the properties and structure of which can be studied by the electron paramagnetic resonance method. [JPRS]

SUB CODE: 07 / SUBM DATE: 21Jun65 / ORIG REF: 006 / OTH REF: 004

Card 2/2 inc

ACC NR: AP7012406

SOURCE CODE: UR/0020/66/170/005/1124/1125

AUTHOR: Valitova, F. G.; Ryzhmanov, Yu. M.

ORG: Kazan Physicotechnical Institute, AN SSSR (Kazanskij fiziko-
tekhnicheskij institut AN SSSR); Institute of Organic and Physical Chemistry
im. A. Ye. Arbuzov, AN SSSR (Institut organicheskoy i fizicheskoy khimii
AN SSSR)

TITLE: Electron paramagnetic resonance in solutions of some free radicals in
the thiophosphonehydrazyl series

SOURCE: AN SSSR. Doklady, v. 170, no. 5, 1966, 1124-1125

TOPIC TAGS: electron paramagnetic resonance, hydrazine compound

SUB CODE: 07

ABSTRACT: The authors studied the isotropic hyperfine structure of the elec-
tron paramagnetic resonance spectra of synthesized diphenylthiophosphonehy-
drazines:



Card 1/2

UDC: 541.67
0932.7336

ACC NR: AP7012406

Solutions of these hydrazines were prepared in chloroform and oxidized with PbO_2 in previously evacuated ampules. The measurements were made on an RE-1301 spectrometer at room temperature. A five-component hyperfine spectrum was observed for all radicals, which may be explained on the basis of interaction between an unpaired electron and two nuclei of nitrogen $\text{N}^{14}(I=1)$

in hydrazine. The constant A in the term of the Hamiltonian $A\vec{I}\vec{S}$ which describes electron-nuclear contact interaction is determined and the results are tabulated. It was experimentally found that substituents have an insignificant effect on the hyperfine spectra. The experimental data show a reduction in the sum of constants A_1+A_2 to little more than half the value observed in DPPG and its derivatives. A probable reason for this decrease is reduction in the admixture of the s -state to the wave function of the unpaired electron for a b -atom of nitrogen. Ratios of $A_1/A_2=1$ were observed for the phenyl, oxyphenyl and oxyethyl radicals which indicates complete conjugation of p_π -orbitals for $\text{N}_\alpha-\text{N}_\beta$. This assumption is confirmed by the planar distribution of all σ -orbitals in $\text{N}_\alpha-\text{N}_\beta$. This article was presented by

Academician D. A. Arbuzov on 31 January 1966. The authors thank B. M. KOZYREV and Yu. V. YABLOKOV for participating in discussion of the results. Orig. art. has: 2 formulas and 1 table. [JPRS: 40,422]

3/2

IVANOV, B.Ye.; VALITOVA, L.A.

Synthesis of substituted benzylphosphinic acids. Izv. AN SSSR.
Otd.khim.nauk no.6:1049-1052 Je '63. (MIRA 16:7)

1. Institut organicheskoy khimii AN SSSR, Kazan'.
(Phosphinic acid) (Rearrangements (Chemistry))

1970-1971
1972-1973
1974-1975
1976-1977
1978-1979
1980-1981
1982-1983
1984-1985
1986-1987
1988-1989
1990-1991
1992-1993
1994-1995
1996-1997
1998-1999
2000-2001
2002-2003
2004-2005
2006-2007
2008-2009
2010-2011
2012-2013
2014-2015
2016-2017
2018-2019
2020-2021
2022-2023
2024-2025
2026-2027
2028-2029
2030-2031
2032-2033
2034-2035
2036-2037
2038-2039
2040-2041
2042-2043
2044-2045
2046-2047
2048-2049
2050-2051
2052-2053
2054-2055
2056-2057
2058-2059
2060-2061
2062-2063
2064-2065
2066-2067
2068-2069
2070-2071
2072-2073
2074-2075
2076-2077
2078-2079
2080-2081
2082-2083
2084-2085
2086-2087
2088-2089
2090-2091
2092-2093
2094-2095
2096-2097
2098-2099
2100-2101
2102-2103
2104-2105
2106-2107
2108-2109
2110-2111
2112-2113
2114-2115
2116-2117
2118-2119
2120-2121
2122-2123
2124-2125
2126-2127
2128-2129
2130-2131
2132-2133
2134-2135
2136-2137
2138-2139
2140-2141
2142-2143
2144-2145
2146-2147
2148-2149
2150-2151
2152-2153
2154-2155
2156-2157
2158-2159
2160-2161
2162-2163
2164-2165
2166-2167
2168-2169
2170-2171
2172-2173
2174-2175
2176-2177
2178-2179
2180-2181
2182-2183
2184-2185
2186-2187
2188-2189
2190-2191
2192-2193
2194-2195
2196-2197
2198-2199
2200-2201
2202-2203
2204-2205
2206-2207
2208-2209
2210-2211
2212-2213
2214-2215
2216-2217
2218-2219
2220-2221
2222-2223
2224-2225
2226-2227
2228-2229
2230-2231
2232-2233
2234-2235
2236-2237
2238-2239
2240-2241
2242-2243
2244-2245
2246-2247
2248-2249
2250-2251
2252-2253
2254-2255
2256-2257
2258-2259
2260-2261
2262-2263
2264-2265
2266-2267
2268-2269
2270-2271
2272-2273
2274-2275
2276-2277
2278-2279
2280-2281
2282-2283
2284-2285
2286-2287
2288-2289
2290-2291
2292-2293
2294-2295
2296-2297
2298-2299
2300-2301
2302-2303
2304-2305
2306-2307
2308-2309
2310-2311
2312-2313
2314-2315
2316-2317
2318-2319
2320-2321
2322-2323
2324-2325
2326-2327
2328-2329
2330-2331
2332-2333
2334-2335
2336-2337
2338-2339
2340-2341
2342-2343
2344-2345
2346-2347
2348-2349
2350-2351
2352-2353
2354-2355
2356-2357
2358-2359
2360-2361
2362-2363
2364-2365
2366-2367
2368-2369
2370-2371
2372-2373
2374-2375
2376-2377
2378-2379
2380-2381
2382-2383
2384-2385
2386-2387
2388-2389
2390-2391
2392-2393
2394-2395
2396-2397
2398-2399
2400-2401
2402-2403
2404-2405
2406-2407
2408-2409
2410-2411
2412-2413
2414-2415
2416-2417
2418-2419
2420-2421
2422-2423
2424-2425
2426-2427
2428-2429
2430-2431
2432-2433
2434-2435
2436-2437
2438-2439
2440-2441
2442-2443
2444-2445
2446-2447
2448-2449
2450-2451
2452-2453
2454-2455
2456-2457
2458-2459
2460-2461
2462-2463
2464-2465
2466-2467
2468-2469
2470-2471
2472-2473
2474-2475
2476-2477
2478-2479
2480-2481
2482-2483
2484-2485
2486-2487
2488-2489
2490-2491
2492-2493
2494-2495
2496-2497
2498-2499
2500-2501
2502-2503
2504-2505
2506-2507
2508-2509
2510-2511
2512-2513
2514-2515
2516-2517
2518-2519
2520-2521
2522-2523
2524-2525
2526-2527
2528-2529
2530-2531
2532-2533
2534-2535
2536-2537
2538-2539
2540-2541
2542-2543
2544-2545
2546-2547
2548-2549
2550-2551
2552-2553
2554-2555
2556-2557
2558-2559
2560-2561
2562-2563
2564-2565
2566-2567
2568-2569
2570-2571
2572-2573
2574-2575
2576-2577
2578-2579
2580-2581
2582-2583
2584-2585
2586-2587
2588-2589
2590-2591
2592-2593
2594-2595
2596-2597
2598-2599
2600-2601
2602-2603
2604-2605
2606-2607
2608-2609
2610-2611
2612-2613
2614-2615
2616-2617
2618-2619
2620-2621
2622-2623
2624-2625
2626-2627
2628-2629
2630-2631
2632-2633
2634-2635
2636-2637
2638-2639
2640-2641
2642-2643
2644-2645
2646-2647
2648-2649
2650-2651
2652-2653
2654-2655
2656-2657
2658-2659
2660-2661
2662-2663
2664-2665
2666-2667
2668-2669
2670-2671
2672-2673
2674-2675
2676-2677
2678-2679
2680-2681
2682-2683
2684-2685
2686-2687
2688-2689
2690-2691
2692-2693
2694-2695
2696-2697
2698-2699
2700-2701
2702-2703
2704-2705
2706-2707
2708-2709
2710-2711
2712-2713
2714-2715
2716-2717
2718-2719
2720-2721
2722-2723
2724-2725
2726-2727
2728-2729
2730-2731
2732-2733
2734-2735
2736-2737
2738-2739
2740-2741
2742-2743
2744-2745
2746-2747
2748-2749
2750-2751
2752-2753
2754-2755
2756-2757
2758-2759
2760-2761
2762-2763
2764-2765
2766-2767
2768-2769
2770-2771
2772-2773
2774-2775
2776-2777
2778-2779
2780-2781
2782-2783
2784-2785
2786-2787
2788-2789
2790-2791
2792-2793
2794-2795
2796-2797
2798-2799
2800-2801
2802-2803
2804-2805
2806-2807
2808-2809
2810-2811
2812-2813
2814-2815
2816-2817
2818-2819
2820-2821
2822-2823
2824-2825
2826-2827
2828-2829
2830-2831
2832-2833
2834-2835
2836-2837
2838-2839
2840-2841
2842-2843
2844-2845
2846-2847
2848-2849
2850-2851
2852-2853
2854-2855
2856-2857
2858-2859
2860-2861
2862-2863
2864-2865
2866-2867
2868-2869
2870-2871
2872-2873
2874-2875
2876-2877
2878-2879
2880-2881
2882-2883
2884-2885
2886-2887
2888-2889
2890-2891
2892-2893
2894-2895
2896-2897
2898-2899
2900-2901
2902-2903
2904-2905
2906-2907
2908-2909
2910-2911
2912-2913
2914-2915
2916-2917
2918-2919
2920-2921
2922-2923
2924-2925
2926-2927
2928-2929
2930-2931
2932-2933
2934-2935
2936-2937
2938-2939
2940-2941
2942-2943
2944-2945
2946-2947
2948-2949
2950-2951
2952-2953
2954-2955
2956-2957
2958-2959
2960-2961
2962-2963
2964-2965
2966-2967
2968-2969
2970-2971
2972-2973
2974-2975
2976-2977
2978-2979
2980-2981
2982-2983
2984-2985
2986-2987
2988-2989
2990-2991
2992-2993
2994-2995
2996-2997
2998-2999
3000-3001
3002-3003
3004-3005
3006-3007
3008-3009
3010-3011
3012-3013
3014-3015
3016-3017
3018-3019
3020-3021
3022-3023
3024-3025
3026-3027
3028-3029
3030-3031
3032-3033
3034-3035
3036-3037
3038-3039
3040-3041
3042-3043
3044-3045
3046-3047
3048-3049
3050-3051
3052-3053
3054-3055
3056-3057
3058-3059
3060-3061
3062-3063
3064-3065
3066-3067
3068-3069
3070-3071
3072-3073
3074-3075
3076-3077
3078-3079
3080-3081
3082-3083
3084-3085
3086-3087
3088-3089
3090-3091
3092-3093
3094-3095
3096-3097
3098-3099
3100-3101
3102-3103
3104-3105
3106-3107
3108-3109
3110-3111
3112-3113
3114-3115
3116-3117
3118-3119
3120-3121
3122-3123
3124-3125
3126-3127
3128-3129
3130-3131
3132-3133
3134-3135
3136-3137
3138-3139
3140-3141
3142-3143
3144-3145
3146-3147
3148-3149
3150-3151
3152-3153
3154-3155
3156-3157
3158-3159
3160-3161
3162-3163
3164-3165
3166-3167
3168-3169
3170-3171
3172-3173
3174-3175
3176-3177
3178-3179
3180-3181
3182-3183
3184-3185
3186-3187
3188-3189
3190-3191
3192-3193
3194-3195
3196-3197
3198-3199
3200-3201
3202-3203
3204-3205
3206-3207
3208-3209
3210-3211
3212-3213
3214-3215
3216-3217
3218-3219
3220-3221
3222-3223
3224-3225
3226-3227
3228-3229
3230-3231
3232-3233
3234-3235
3236-3237
3238-3239
3240-3241
3242-3243
3244-3245
3246-3247
3248-3249
3250-3251
3252-3253
3254-3255
3256-3257
3258-3259
3260-3261
3262-3263
3264-3265
3266-3267
3268-3269
3270-3271
3272-3273
3274-3275
3276-3277
3278-3279
3280-3281
3282-3283
3284-3285
3286-3287
3288-3289
3290-3291
3292-3293
3294-3295
3296-3297
3298-3299
3300-3301
3302-3303
3304-3305
3306-3307
3308-3309
3310-3311
3312-3313
3314-3315
3316-3317
3318-3319
3320-3321
3322-3323
3324-3325
3326-3327
3328-3329
3330-3331
3332-3333
3334-3335
3336-3337
3338-3339
3340-3341
3342-3343
3344-3345
3346-3347
3348-3349
3350-3351
3352-3353
3354-3355
3356-3357
3358-3359
3360-3361
3362-3363
3364-3365
3366-3367
3368-3369
3370-3371
3372-3373
3374-3375
3376-3377
3378-3379
3380-3381
3382-3383
3384-3385
3386-3387
3388-3389
3390-3391
3392-3393
3394-3395
3396-3397
3398-3399
3400-3401
3402-3403
3404-3405
3406-3407
3408-3409
3410-3411
3412-3413
3414-3415
3416-3417
3418-3419
3420-3421
3422-3423
3424-3425
3426-3427
3428-3429
3430-3431
3432-3433
3434-3435
3436-3437
3438-3439
3440-3441
3442-3443
3444-3445
3446-3447
3448-3449
3450-3451
3452-3453
3454-3455
3456-3457
3458-3459
3460-3461
3462-3463
3464-3465
3466-3467
3468-3469
3470-3471
3472-3473
3474-3475
3476-3477
3478-3479
3480-3481
3482-3483
3484-3485
3486-3487
3488-3489
3490-3491
3492-3493
3494-3495
3496-3497
3498-3499
3500-3501
3502-3503
3504-3505
3506-3507
3508-3509
3510-3511
3512-3513
3514-3515
3516-3517
3518-3519
3520-3521
3522-3523
3524-3525
3526-3527
3528-3529
3530-3531
3532-3533
3534-3535
3536-3537
3538-3539
3540-3541
3542-3543
3544-3545
3546-3547
3548-3549
3550-3551
3552-3553
3554-3555
3556-3557
3558-3559
3560-3561
3562-3563
3564-3565
3566-3567
3568-3569
3570-3571
3572-3573
3574-3575
3576-3577
3578-3579
3580-3581
3582-3583
3584-3585
3586-3587
3588-3589
3590-3591
3592-3593
3594-3595
3596-3597
3598-3599
3600-3601
3602-3603
3604-3605
3606-3607
3608-3609
3610-3611
3612-3613
3614-3615
3616-3617
3618-3619
3620-3621
3622-3623
3624-3625
3626-3627
3628-3629
3630-3631
3632-3633
3634-3635
3636-3637
3638-3639
3640-3641
3642-3643
3644-3645
3646-3647
3648-3649
3650-3651
3652-3653
3654-3655
3656-3657
3658-3659
3660-3661
3662-3663
3664-3665
3666-3667
3668-3669
3670-3671
3672-3673
3674-3675
3676-3677
3678-3679
3680-3681
3682-3683
3684-3685
3686-3687
3688-3689
3690-3691
3692-3693
3694-3695
3696-3697
3698-3699
3700-3701
3702-3703
3704-3705
3706-3707
3708-3709
3710-3711
3712-3713
3714-3715
3716-3717
3718-3719
3720-3721
3722-3723
3724-3725
3726-3727
3728-3729
3730-3731
3732-3733
3734-3735
3736-3737
3738-3739
3740-3741
3742-3743
3744-3745
3746-3747
3748-3749
3750-3751
3752-3753
3754-3755
3756-3757
3758-3759
3760-3761
3762-3763
3764-3765
3766-3767
3768-3769
3770-3771
3772-3773
3774-3775
3776-3777
3778-3779
3780-3781
3782-3783
3784-3785
3786-3787
3788-3789
3790-3791
3792-3793
3794-3795
3796-3797
3798-3799
3800-3801
3802-3803
3804-3805
3806-3807
3808-3809
3810-3811
3812-3813
3814-3815
3816-3817
3818-3819
3820-3821
3822-3823
3824-3825
3826-3827
3828-3829
3830-3831
3832-3833
3834-3835
3836-3837
3838-3839
3840-3841
3842-3843
3844-3845
3846-3847
3848-3849
3850-3851
3852-3853
3854-3855
3856-3857
3858-3859
3860-3861
3862-3863
3864-3865
3866-3867
3868-3869
3870-3871
3872-3873
3874-3875
3876-3877
3878-3879
3880-3881
3882-3883
3884-3885
3886-3887
3888-3889
3890-3891
3892-3893
3894-3895
3896-3897
3898-3899
3900-3901
3902-3903
3904-3905
3906-3907
3908-3909
3910-3911
3912-3913
3914-3915
3916-3917
3918-3919
3920-3921
3922-3923
3924-3925
3926-3927
3928-3929
3930-3931
3932-3933
3934-3935
3936-3937
3938-3939
3940-3941
3942-3943
3944-3945
3946-3947
3948-3949
3950-3951
3952-3953
3954-3955
3956-3957
3958-3959
3960-3961
3962-3963
3964-3965
3966-3967
3968-3969
3970-3971
3972-3973
3974-3975
3976-3977
3978-3979
3980-3981
3982-3983
3984-3985
3986-3987
3988-3989
3990-3991
3992-3993
3994-3995
3996-3997
3998-3999
4000-4001
4002-4003
4004-4005
4006-4007
4008-4009
4010-4011
4012-4013
4014-4015
4016-4017
4018-4019
4020-4021
4022-4023
4024-4025
4026-4027
4028-4029
4030-4031
4032-4033
4034-4035
4036-4037
4038-4039
4040-4041
4042-4043
4044-4045
4046-4047
4048-4049
4050-4051
4052-4053
4054-4055
4056-4057
4058-4059
4060-4061
4062-4063
4064-4065
4066-4067
4068-4069
4070-4071
4072-4073
4074-4075
4076-4077
4078-4079
4080-4081
4082-4083
4084-4085
4086-4087
4088-4089
4090-4091
4092-4093
4094-4095
4096-4097
4098-4099
4100-4101
4102-4103
4104-4105
4106-4107
4108-4109
4110-4111
4112-4113
4114-4115
4116-4117
4118-4119
4120-4121
4122-4123
4124-4125
4126-4127
4128-4129
4130-4131
4132-4133
4134-4135
4136-4137
4138-4139
4140-4141
4142-4143
4144-4145
4146-4147
4148-4149
4150-4151
4152-4153
4154-4155
4156-4157
4158-4159
4160-4161
4162-4163
4164-4165
4166-4167
4168-4169
4170-4171
4172-4173
4174-4175
4176-4177
4178-4179
4180-4181
4182-4183
4184-4185
4186-4187
4188-4189
4190-4191
4192-4193
4194-4195
4196-

acid or hydroxylbenzyl acetate, and the ester of the polytransacetylation of the diethyl ester of 4,4'-oxybis(benzophenonic acid)

SUBMITTED: 1965

DATE: 1965

NO. ONE

NO. TWO

NO.

NO.

Card

VALITOVA, M. S. Cand Med Sci -- (diss) "On Age ^{related} Changes in the Content of Myoglobin, Hemoglobin, and Iron in the Organism of Dogs." Alma-Ata, 1957. 10 pp 21 cm. (Kazakh ~~XXXXXXXXXX~~ Medical Inst im V. M. Molotov), 300 copies (KL, 26-57, 112)

- 112 -

VALITOVA, M.S.

VALITOVA, M.S.

Changes with age in the amount of ~~myoglobin~~, hemoglobin, and iron
in the organism of dogs [with summary in English]. Ukr.biokhim.
zhur. 29 no.4:419-427 '57. (MIRA 11:1)

1. Kafedra biokhimii Kazakhskogo meditsinskogo instituta, Alma-Ata.
(MYOGLOBIN) (HEMOGLOBIN) (IRON IN THE BODY)

VALITOVA, M.S., DYKOVA, A.L., KEROVSKAYA, N.I., FANTUA, Z.A.,
RESHETNIKOVA, M.I., SULAYEVA, L.S., UTESHEV, A.B., VERBOLOBICH, P.A.,
POLOSUKHINA, T. YA., KAIPOVA, Z.M., (USSR)

"Special Aspects of the Metabolism of Some Substances in
Radiation Disease in Dogs."

Report presented at the 5th Int'l. Biochemistry Congress, Moscow,
10-16 Aug 1961.

47,2400
Z/011/62/019/011/001/003
E073/E535

AUTHORS: Polosukhina T. V., Valitova M.S. et al.

TITLE: Influence of X-ray irradiation on the metabolism of lipides in the liver of a dog.

PERIODICAL: Chemie a chemická technologie. Přehled technické a hospodářské literatury, v.19, no.11, 1962, 496 abstract Ch 62 6690 (Vopr. med. Khim. v.8, no.2.1962, 192-199)

TEXT: The serum of venous blood showed an increased concentration of neutral fats, phosphor lipides and cholesterol the third day after a single irradiation of the dog with an X-ray dose (500 roentgen). The increase in the concentration of ketones in the blood in the case of radiation sickness is caused by the formation of ketones in the liver. The content of glycogens in the liver tissue decreased, whilst the content of neutral fats and cholesterol increased.
3 tables, 46 references.

[Abstracter's note: Complete translation.]

Card 1/1

1ST AND 2ND ORDERS										3RD AND 4TH ORDERS									
PROCESSES AND PROPERTIES INDEX																			
<p>20</p> <p>The Nishmetarbagat manganese deposit. - I. M. Khalilovskaya. <i>Razvedka Nedr</i> 1957, No. 3, 13-14. - The ore contains MnO₂ 11.37-17.85, Mn 7.10-11.11, SiO₂ 66.77-61.00, Al₂O₃ 13.73-16.21, Fe₂O₃ 1.43-4.30, S 0.02, CaO 1.11-0.61 and MgO 0.03-1.10%.</p> <p>A. A. Bozhilovsk</p>																			
<p>ASB-5LA METALLURGICAL LITERATURE CLASSIFICATION</p>																			
<p>10000 02 10000 02 10000 02 10000 02 10000 02 10000 02 10000 02 10000 02 10000 02 10000 02</p>																			

VALITSKIY, I.V.

VALITSKIY, I.V., redaktor; IOANNESYAN, R.A., redaktor; KOPYSITSKIY, P.I.,
redaktor; GAUMIN, I.M., redaktor; TROFIMOV, A.V., tekhnicheskii
redaktor.

[Improving the quality of cutting bits; collection of articles]
Povyshenie kachestva izgotovleniya sharoshechnykh dolot; sbornik
statei. Moskva, Gos.nauchno-tekhn.izd-vo neftianoi i gorno-toplivnoi
lit-ry, 1954. 93 p. (MIRA 8:4)

1. Moscow. TSentral'nyy nauchno-issledovatel'skiy institut mekhanizatsii i organizatsii truda v neftyanoy promyshlennosti. Byuro tekhniko-ekonomicheskoy informatsii.
(Boring machinery)

VALITSKIY, V., inzh.

Tuning of the TsRS radio station without instruments. Pozh.
delo 8 no.9:24 S '62. (MIRA 16:11)

VALITSKIY, V. P. AND V. O. ARUTUNOV

Elektroizmeritel'nye pribory; ustroistvo, montazh i obsluzhivanie. Izd. 2.
Leningrad, Gosenergoizdat, 1949. 199 p. illus.

Electric meters; working principles, adjustment and maintenance.

DLC: TK393.A7 1949

SO: Manufacturing and Mechanical Engineering in the Soviet Union, Library of
Congress, 1953.

VALITSKIY, V.P. and ARUTYUNOV, V.O.

"Electrical Measuring Instruments". (Elektroizmeritel'nyye pribory),
Gosenergoizdat, Leningrad/Moscow, 1949, 199 pp, 7 rubles 50 kopeks,

SO: W-14151 11 Oct 1950.

"APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001858510004-9

APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001858510004-9"

"APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001858510004-9

APPROVED FOR RELEASE: 08/31/2001

CIA-RDP86-00513R001858510004-9"

ACC NR: AP6032848
 SOURCE CODE: UR/0020/66/170/003/0540/0543
 AUTHOR: Belyakov, L. V.; Valitskiy, V. P.; Zlatin, N. A.; Mochalov, S. M.
 ORG: Physical-Technical Institute im. A. F. Ioffe, Academy of Sciences SSSR (Fiziko-
 tekhnicheskiy institut Akademii nauk SSSR)
 TITLE: The melting of lead in a shock wave
 SOURCE: AN SSSR. Doklady, v. 170, no. 3, 1966, 540-543
 TOPIC TAGS: shock wave, x ray photography, high speed camera, pressure distribution,
 specific volume, thermodynamic analysis
 ABSTRACT: A study was made of adiabatic heating of lead to the fusion point during im-
 pact shock loading. Thermodynamic analysis of melting in a shock wave is presented
 and schematic drawings are given of pressure as a function of specific volume and dis-
 tance. Thermodynamic equations are given for the specific work done by pressure to
 $\alpha\lambda$, where λ is the specific heat of fusion and α is a coefficient which depends on the
 shock pressure. It is shown that in an entropy increase and a change
 in pressure distribution. These analytical results were checked by shock wave experi-
 ments on lead, in which high speed x-ray photographs were taken of the fractured ends
 of lead sheets. Impact velocities ranged from 1085 to 1570 m/sec. A sharp change in
 fracture appearance occurred at an impact velocity of 1250-1300 m/sec; this coincided

ACC NR: AP6032848
 SOURCE CODE: UR/0020/66/170/003/0540/0543

AUTHOR: Belyakov, L. V.; Valitskiy, V. P.; Zlatin, N. A.; Mochalov, S. M.

ORG: Physical-Technical Institute im. A. F. Ioffe, Academy of Sciences SSSR (Fiziko-
 tekhnicheskiy institut Akademii nauk SSSR)

TITLE: The melting of lead in a shock wave

SOURCE: AN SSSR. Doklady, v. 170, no. 3, 1966, 540-543

TOPIC TAGS: shock wave, x ray photography, high speed camera, pressure distribution,
 specific volume, thermodynamic analysis

ABSTRACT: A study was made of adiabatic heating of lead to the fusion point during im-
 pact shock loading. Thermodynamic analysis of melting in a shock wave is presented
 and schematic drawings are given of pressure as a function of specific volume and dis-
 tance. Thermodynamic equations are given for the specific work done by pressure to
 $\alpha\lambda$, where λ is the specific heat of fusion and α is a coefficient which depends on the
 shock pressure. It is shown that in an entropy increase and a change
 in pressure distribution. These analytical results were checked by shock wave experi-
 ments on lead, in which high speed x-ray photographs were taken of the fractured ends
 of lead sheets. Impact velocities ranged from 1085 to 1570 m/sec. A sharp change in
 fracture appearance occurred at an impact velocity of 1250-1300 m/sec; this coincided

Card 1/2

UDC: 531.66.001.11

L 07443-067

ACC NR: AP6032848

3
with a mass velocity of 700 m/sec. It is known that melting of lead occurs in a shock wave when the mass velocity becomes 650-700 m/sec. This velocity corresponded to a pressure of $230-250 \times 10^3$ atm and to a 22-23% change in specific volume. X-ray photographs are also shown of fracture in 1 mm thick lead sheets at an impact velocity of 1340 m/sec, during time intervals of 15, 30, 45, and 54 μ sec. These tests show that the difference between the speed of the split flange (initial fracture condition) and the residual mass of the "whiskers", (final fracture condition) was 3%, verifying the specific work equation. Sheet thicknesses ranging from 0.5 to 3 mm were tested 15 μ sec after the moment of fracture at 1340 m/sec. Some of the sheets were covered with 0.05 mm thick aluminum foil during testing. The use of the foil changed the spacings of cleavage "whiskers". These experiments confirmed that the originally postulated position of shock adiabates of lead in the solid and two-phase conditions was correct. A calculation of the relaxation time from the data gave 3×10^{-7} sec. Presented by Academician B. P. Konstantinov on 13 December 1965. Orig. art. has: 4 figures, 2 formulas.

SUB CODE: 11/ SUBM DATE: 27Nov65/ ORIG REF: 008/ OTH REF: 001

Card 2/2 *ala*

L 07367-67 EWT(d)/EWT(l)/EWT(m)/EWP(c)/EWP(w)/EWP(t)/ETI IJP(c) JD/EM/JH
ACC NR: AP6033425 SOURCE CODE: UR/0057/66/036/010/1375/1882

AUTHOR: Belyakov, L. V.; Valitskiy, V. P.; Zlatin, N. A.

ORG: Physicotechnical Institute im. A. F. Ioffe, AN SSSR, Leningrad (Fiziko-
tekhnicheskii institut AN SSSR)

TITLE: Thermal ^{test} effects accompanying an impact on a metal half-space

SOURCE: Zhurnal tekhnicheskoy fiziki, v. 36, no. 10, 1966, 1875-1882

TOPIC TAGS: impact, impact ^{test} effect, ~~impact~~ thermal effect, impact effect modeling,
~~metal test~~

ABSTRACT: The article discusses experiments aimed at expanding the range of applicability of criteria to parallel the effects between low-speed impact of one pair of materials to similar effects for a pair of different materials at a higher, experimentally unattainable, speed. The upper limit of the interval for which the modeling curve will yield correct results is discussed at some length. The concept of "threshold speed" is introduced. Threshold speed corresponds to the discontinuity of the modeling curve caused by melting of the metals in question and is estimated at about 0.7 to 1 of the velocity of sound in the given metal. If the threshold speed is correctly determined, the modeling curve should yield accurate data on impact results for speeds at least 3 to 3.5 times higher than the experimental. Experiments were conducted in which blocks of lead, tin, and cadmium were impacted by aluminum disks 4 mm thick and 15 mm in diameter at speeds up to 24 km/sec.

Card 1/2

UDC: 531.66.001.11

L 07367-67

ACC NR: AP6033425

The results were in good agreement with calculated threshold speeds. A marked difference was observed in the cavities formed at impact speeds of about 1 km/sec and those obtained at about 1.7 to 1.8 km/sec, the latter showing conical deepening with evidence of fusing of the target. A special high-speed x-ray investigation of the process at impact speeds up to 2.4 km/sec confirmed the ejection of molten material at speeds of 1.7 to 1.8 km/sec and higher. Further confirmation of the threshold speed magnitudes was obtained in a special series of x-ray tests in which a copper cylinder was made to hit thin (about 0.05 of the diameter of the cylinder) sheets of lead, tin, and cadmium. The threshold speeds for metals with higher melting point were calculated on the assumption that the heating up of the target by the impact is a function of the mass speed developed by the shock load and obeys the same law for all metals. The results of calculations for a number of metals confirm the assumption and agree with experimental data from various sources. Reference is made to the experiments of A. C. Charters (Sci. Amer. v. 203, no. 4, 1960, 128), whose results could be extrapolated for impact speeds of 30 to 50 km/sec. Orig. art. has: 6 figures.

SUB CODE: 20/ SUBM DATE: 18Oct65/ ORIG REF: 012/ OTH REF: 007/ ATD PRESS: 5101

Card 2/2 afe

AUTHOR: FISHMAN, K.M., VALITSKIY, Yu.N.

20-6-5/47

TITLE: On the Applicability of the Fredholm Theory to Some Linear Topological Spaces (O primenimosti teorii Fredgol'ma k nekotorym lineynym topologicheskim prostranstvam)

PERIODICAL: Doklady Akademii Nauk ~~SSSR~~, 1957, Vol. 117, Nr 6, pp 943-946 (USSR)

ABSTRACT: Let B_r be a real or complex Banach space depending on a parameter r ($\alpha < r \leq \beta$) and satisfying certain further conditions. In the set $\mathcal{O}_r = \bigcap_{r' < r} B_{r'}$ ($\alpha < r \leq \beta$) let the addition and convergence be defined such that \mathcal{O}_r becomes a complete linear topological space. Let A be a linear operator which maps B_r into B_r and which is independent of r . In every \mathcal{O}_r , A induces a linear continuous operator not depending on r . The author shows that if the Fredholm theory can be applied to A in every B_r , then also in the spaces \mathcal{O}_r all Fredholm theorems are valid.

Card 1/2

4 Soviet references are quoted.

' On the Applicability of the Fredholm Theory to Some Linear
Topological Spaces 20-6-5/47

ASSOCIATION. Chernovtsy State University (Chernovitskiy gosudarstvennyy
universitet)
PRESENTED: By V.I.Smirnov, Academician, 21 June 1957
SUBMITTED: 20 June 1957
AVAILABLE: Library of Congress

Card 2/2

SOV/21-59-3-3/27

16(1)

AUTHOR:

Valitskiy, Yu.N.

TITLE:

Functions, Analytical with Respect to Some Integro-Differential Operators, and Their Application
(Funktsii analiticheskiye otnositel'no nekotorykh integro-differentsial'nykh operatorov i ikh primeneniye)

PERIODICAL:

Dopovidi Akademii nauk Ukrain's'koi RSR, 1959, Nr 3, pp 237-240 (USSR)

ABSTRACT:

The object of this article is to accomplish an approach generalizing a theory of functions analytical with respect to a conventional operator of the n -th order, and proving that two such operators are equivalent, so as to make it fit for use in cases with one type of integro-differential operators. Step by step, considering and proving eight theorems, the author established the facts which hold in the theory of ordinary analytical functions and are fit to serve for a number of functions that can be expanded into series. The terminology employed

Card 1/2

SOV/21-59-3-3/27

Functions, Analytical with Respect to Some Integro-Differential Operators, and Their Application

is standard mathematical. There are 6 Soviet references.

ASSOCIATION: Chernovitskiy gosudarstvennyy universitet (Chernovtsy State University)

PRESENTED: November 29, 1958, by B.V. Gnedenko, Member of the AS UkrSSR

Card 2/2

VALITSKIY, Yu.N.; BREZOVSKIY, N.I.

First Abelian theorem for certain functional series. Vop. mat. fiz.
i teor. funk. no.1:12-17 '64. (MIRA 18:2)

L 16377-67 EWT(1) SCTE DD
ACC NM AP6035944 (N) SOURCE CODE: UR/0413/66/000/020/0214/0214 35

INVENTOR: Grebennikov, N. P.; Valiulin, A. Z.

ORG: none

TITLE: Training device for swimmers. Class 77, No. 187577

SOURCE: Izobreteniya, promyshlennyye obraztsy, tovarnyye znaki, no. 20, 1966,
214

TOPIC TAGS: training equipment, training, potentiometer, liquid flow, hydraulic
pump

ABSTRACT: An Author Certificate has been issued for a training device for swimmers. It is a basin with a closed running-water channel containing a hydraulic pump, which controls the flow rate in the basin. The stream's rate of flow is regulated by a flow sensor attached to the swimmer and consisting of a small cable joined to a coil with a spring which activates the slide bar of a potentiometer regulating the rpm of the pump's motor. Flow distribution grids insure an even rate of water flow through its cross section and are situated at the entrance and exit of the basin. Orig. art. has: 1 figure. [Translation] [N-67-2]


Card 1/2

UDC: 685.734

L 10877-67

ACC NR: AP6035944

Fig. 1. Training device for swimmers.
1—Basin; 2—water flow channel;
3—hydraulic pump; 4—flow sensor;
5—flow distribution grids



SUB CODE: 13, 06/ SUBM DATE: 15Mar65/

Card 2/2

VALIULINA A.Z.

VALIULINA, A.Z.

TOMASHOV, N.D.; KIPARISOV, G.N.; VALIULINA, A.Z.; KOROTKOVA, K.S.

Apparatus for obtaining polarization curves. Trudy Inst. Fiz.Khim.,
Akad. Nauk S.S.S.R. 3. Issledovaniya Korrozii Metal. No.2, 74-5 '51.
(CA 47 no.16:7831 '53) (MLRA 4:10)

VALIULINA, A. Z.

USSR/Chemistry - Corrosion

Mar 52

"Ozone as a Cathodic Depolarizer in Processes of Metal Corrosion," N. D. Tomashov,
A. Z. Valiulina, Inst of Phys Chem, Moscow, Acad Sci USSR

"Zhur Fiz Khim" Vol XXVI, No 3, pp 417-424

In a weakly alk soln ($pH=9.2$) ozone is not reduced directly: decomp at the cathode, it produces supersatn with oxygen and increases the limiting diffusion current. In acidic electrolytes, at much more pos potentials than the potential of oxygen ionization, ozone is directly reduced under assimilation of 2 electrons. The limiting diffusion current is higher in an acidic than alk medium, because ozone is more stable in acidic soln. The authors' assumption that ozone is a corrosion accelerator because of its cathodic depolarizing effect has been confirmed experimentally. In the expts described, Cu, Fe, and Al were as cathodic materials. The problem of corrosion caused by ozone is of importance at high voltage installations.

PA 213T34

VALIULLINA, F.G.

Electrophoresis using the vitreous body in treating a trachomatous pannus and a persistent turbidity of the cornea. Nauch. trudy Kaz. gos. med. inst. 14:383-384 '64. (MIRA 18:9)

1. Kafedra glaznykh bolezney (zav. - doktor med. nauk A.P. Nesterov) Kazanskogo meditsinskogo in-ta.

Valiulina, F.M.

USSR /Chemical Technology. Chemical Products
and Their Application

I-16

Treatment of natural gases and petroleum.
Motor fuels. Lubricants

Abs Jour: Referat Zhur - Khimiya, No 9, 1957, 31897

Author : Rumyantseva Z. A., Valiulina F.M., Chayko V.P.

Inst : Academy of Sciences Tadzhik SSR

Title : Chemical Nature of High Molecular Components of
High Sulfur Petroleum. Communication I. Analysis
of the Components of Petroleum of the Kzyl-Tumshuk
Deposit

Orig Pub: Tr. AN TadzhSSR, 1955, 41, 59-68

Abstract: By methods of GrozNII and adsorption chromato-
graphy on silica gel, a study was made of the

Card 1/4

USSR /Chemical Technology. Chemical Products
and Their Application

I-16

Treatment of natural gases and petroleum.
Motor fuels. Lubricants

Abs Jour: Referat Zhur - Khimiya, No 9, 1957, 31897

groupwise hydrocarbon composition of ligroin-kerosene and oil fractions, and also of the hydrocarbon portion, from which gasoline had been removed, of high-sulfur, Tadzhik petroleum of the Kzyl-Tumshuk deposit, containing about 5% S and 42% of tarry substances. On the basis of fraction yields of the adsorption analysis and their refractive indices, graphs were plotted which show the quantitative distribution of different classes of hydrocarbons in the products under study, and on the basis of determinations of the molecular weight values and elemental composition,

Card 2/4

USSR /Chemical Technology. Chemical Products
and Their Application

I-16

Treatment of natural gases and petroleum.
Motor fuels. Lubricants.

Abs Jour: Referat Zhur - Khimiya, No 9, 1957, 31897

empirical formulas and formulas of homologous series of the fractions were calculated. It is shown that the principal components of the fractions under study are aromatic and sulfur compounds (amounting together to 50-97%); the content of sulfur compounds increases with rising boiling point of the fractions, and on transition from monocyclic aromatic compounds to bicyclic aromatic compounds, and reaches a maximum in the case of separation of a sulfur concentrate. By a ring analysis of the methane-naphthene portion of the fractions it was shown that paraffin hydrocarbons and chains

Card 3/4

USSR /Chemical Technology. Chemical Products
and Their Application

I-16

Treatment of natural gases and petroleum.
Motor fuels. Lubricants.

Abs Jour: Referat Zhur - Khimiya, No 9, 1957, 31897

predominate, and monocyclic naphthenic hydrocarbons are present in the fractions up to 400° and bicyclic naphthenic hydrocarbons in the 400-500° fraction. The Kzyl-Tumshuk petroleum is a rich natural source of organic sulfur compounds.

Card 4/4

RUMYANTSEVA, Z.A.; VALIULINA, F.M.

Chemical nature of components of high molecular weight of highly sulfurous petroleum. Report no. 2. Analyzing the components of Khandag petroleum. Izv. Otd. est. nauk AN Tadzh. SSR no.16:15-28 '56. (MLRA 10:4)

1. Institut khimii AN Tadzhikskoy SSR.
(Khandag--Petroleum--Analysis)

KARAVAYEV, N.M.; RUMYANTSEVA, Z.A.; VALIULINA, F.M.; BURYAKOVA, E.P.

Semicoking of slightly caking and noncaking coal of the
Fan-Yagnob deposit. Izv. Otd. est. nauk AN Tadzh. SSR
no.3:27-38 '59. (MIRA 15:5)

1. Institut khimii AN Tadzhikskoy SSR.
(Ayni District—Coal—Carbonization)

VALIULLIN, A.V.; GIL'MANOV, I.G.; KHASANOV, Kh.Kh.; KOROL'CHUKA, V.M.,
red.; LODVIKOVA, A.S., red. izd-va; NABIULLINA, R.S., tekhn.
red.

[Fruit culture of the Tatar A.S.S.R.] Sadovodstvo Tatarskoi ASSSR.
Kazan', Tatarskoe knizhnoe izd-vo, 1960. 279 p. (MIRA 14:9)
(Tatar A.S.S.R.—Fruit culture)

AYUPOV, Kh.V., kand. veter. nauk; IVANOVSKIY, S.A., kand. veter. nauk;
SAFIULLIN, G.K.; VALIULLIN, S.M., veterinarnyy vrach;
UPORNIKOV, M.V., veterinarnyy vrach; FROLOV, V.P., zootekhnik

Veterinary helminthological evaluation of the year-round
pen system of keeping sheep. Veterinariia 40 no.6:49-52
Je '63. (MIRA 17:1)

1. Bashkirsкая nauchno-proizvodstvennaya veterinarnaya
laboratoriya (for Frolov). 2. Direktor Miyakinskogo sovkhoza
Bashkirskey ASSR (for Safiullin).

VALIULLINA, F.G., assistant

Tissue therapy for trachomatous pannus. Oft. zhur. 14 no.2:118-121
'59. (MIRA 12:7)

1. Iz kafedry glaznykh bolezney (zav. - dotsent A.S. Veyz)
Kazanskogo meditsinskogo instituta.
(CORNEA--DISEASES) (TISSUE EXTRACTS)

KAMCHATNOV, V.P.; VALIULLINA, F.G.; SAMOYLOVA, A.I.

Study of working conditions and incidence of disease in working with methanol in the "dark shops" of the V.V. Kuibyshev Chemical Factory. Kaz. med. zhur. 41 no.3:78-82 My-Je '60. (MIRA 13:9)

1. Iz kafedr gigiyeny truda (zav. - dotsent V.P.Kamchatnov) i glaznykh bolezney (zav. - dotsent A.S. Veys) Kazanskogo meditsinskogo instituta.

(METHYL ALCOHOL—PHYSIOLOGICAL EFFECT)
(CHEMICAL INDUSTRIES—HYGIENIC ASPECTS)

VALJILLINA, F.G., kand.med.nauk

Microsymptoms in methyl alcohol poisoning. Vest.oft. no.3:62-
64 '61. (MIRA 14:9)

1. Kaf'edra glasnykh bolezney (zav. - dotsent A.S. Voys) Kazan-
skogo meditsinskogo instituta.
(METHANOL—TOXICOLOGY)

RADBIL', O.S., prof.; VALIULLINA, R.K.; KOVALERCHIK, E.I.;
FEREL'SHTEYN, S.B.

Clinical aspects and treatment of chronic nonspecific ulcerative
colitis. Terap.arkh. no.8:11-16 '62. (MIRA 15:12)

1. Iz 2-y kafedry terapii (zav. - prof. O.S. Radbil') Kazanskogo
gosudarstvennogo instituta dlya usovershenstvovaniya vrachey i
2-y infektsionnoy bol'nitsy (glavnyy vrach E.I. Kovalerchik).
(COLITIS)

VALIULLINA, R. I.

RYABUKHIN, G.Ye.; VALIULLINA, R.T.

Some new data concerning the facies and lithology of Jurassic and lower Cretaceous sediments of the Bolsherechye deep hole region in the West Siberian Lowland. Dokl. AN SSSR 110 no.6: 1065-1069 0 '56. (MLRA 10:2)

1. Sverdlovskiy gornyy institut imeni V.V. Vakhrusheva.
Predstavleno akademikom S.I. Mironovym.
(Siberia, Western--Geology, Stratigraphic)

VALIULLINA, R.T.; CHERNOMORSKIY, V.N.

Correlation of the lower Carboniferous terrigenous stratum of
northwestern Bashkiria based on the palynological and ~~mineralogical~~
analysis of clayey rocks. Dokl. AN SSSR 139 no.5:1181-1184
Ag. '61. (MIRA 14:8)

1. Predstavleno akademikom D.V. Nalivkinym.
(Bashkiria—Geology, Stratigraphic)

VALIUS, N.A.

[Photography in darkness and fog] Fotos'saka v temnote i v tume.
Moskvan Gostekhisdat, 1948. (MLRA 9:3)

(Photography)

1. VALTUZHENICH, A.
2. USSR (600)
4. Argentina - Social Conditions
7. Beneath the yoke. Mol. kolkh. 20, no. 2, 1953.

9. Monthly List of Russian Accessions, Library of Congress, May 1953. Unclassified.

USMANOV, Kh.U.; YUL'CHIBAYEV, A.A.; MUKHAMEDZHANOV, R.; GORDIYENKO, A.A.;
PATENKO, A.A.; DORDZHIN, G.S.; VALIYEV, A.

Radiation-induced polymerization of vinyl fluoride. Khim. i fiz.-
khim. prirod. i sint. polim. no.18205-206 '62 (MIRA 18:1)

1. Chlen-korrespondent AN UZSSR (for Usmanov).

ACCESSION NR: AP000471

AUTHOR: Usmanov, Kh. U.; Yul'chibayev, A. A.; Mukhamedzhanov, R.; Gordiyenko, A. A.; Valiyev, A.; Patenko, A. A.; Dordzhin, G. S.

TITLE: Radiation-induced polymerization of vinyl fluoride

SOURCE: Vyssokomolekulyarnyye soedineniya, v. 5, no. 8, 1963, 1277

TOPIC TAGS: fluorocarbon polymer, poly(vinyl fluoride), poly(vinyl chloride), Fluoroplast-4, Teflon polytetrafluoroethylene, radiation-induced polymerization, benzoyl peroxide, radical polymerization

ABSTRACT: The higher heat, chemical, and light resistance of poly(vinyl fluoride) (PVF) as compared to poly(vinyl chloride) and the possibility of substituting PVF in certain cases for Fluoroplast-4 [polytetrafluoroethylene, or Teflon] have prompted a study of the synthesis of PVF by radiation-induced polymerization. The monomer was prepared from pure HF and C_2H_2 , separated from excess HF and C_2H_2 , and irradiated in sealed ampoules with γ -rays from a Co^{60} source at a dose rate of 10^5 rad/sec. Irradiation in the absence of initiators yielded waxy products. In the presence of benzoyl peroxide a yellowish solid product was obtained.

Card 1/2

The PVF yield increased in both cases with an increase in the radiation dose. Acceleration of the polymerization in the presence of the initiator indicates that the polymerization follows the radical mechanism.

ASSOCIATION: none

SUBMITTED: 11Mar63

DATE ACQ: 28Aug63

ENCL: 00

SUB CODE: CH, MA

NO REF SOV: 000

OTHER: 002

Card 2/2

VALIYEV, A.; SAMANOV, Zh.; USMANOV, Kh.

Division and comparison of Cretaceous sediments in the eastern
part of the Barsakel'mes trough. Uzb.geol.zhur. 8 no.3:19-23
'64. (MIRA 18:12)

1. Institut geologii i razrabotki neftyanykh i gazovykh
mestorozhdeniy Gosudarstvennogo geologicheskogo komiteta
SSSR. Submitted Jan. 14, 1964.

VALIYEV, A. A., CAND GEOL-MIN SCI, ^{//} ~~THE~~ EXPERIENCE OF
~~THE~~ PALEOMAGNETIC DISINTEGRATION OF ~~THE~~ CENOZOIC MOLASSES
OF THE CHUST-PAP ANTICLINE. ^{//} LENINGRAD, 1960. (ALL-UNION
PETROLEUM SCI RES GEOL ^{Prospecting} ~~ENRICH~~ INST, ACAD SCI UZSSR. INST OF
GEOL AND ^{Mining} ~~EXPLOITATION~~ OF PETR AND GAS DEPOSITS). (KL, 2-61,
201).

S/169/62/000/010/014/071
D228/D307

AUTHOR: Valiyev, A.A.
TITLE: Magnetic properties of the continental Cenozoic
Molasse of certain Fergana districts
PERIODICAL: Referativnyy zhurnal, Geofizika, no. 10, 1962, 14,
abstract 10A87 (Dokl. AN UzSSR, no. 4, 1960, 13-16
(summary in Uzbek))

TEXT: The susceptibility χ and the remanent magnetization I_n of the Molasse near the Chust-Papskaya Anticline, the Akbel'skaya Ridge, and the Belesenyk Range are briefly reviewed. The magnetic characteristic variation curves for two profiles are given together with the typical magnitudes of χ and I_n for the remaining districts. It is pointed out that the significance of χ and I_n as stratigraphic correlatives is lessened by the fact that they are not sustained, even on nearby profiles having the dissimilar composition of the supply provinces. Only profiles, in which the composition of the products of the supply provinces is identical and the history of tect-

Card 1/2

Magnetic properties ...

S/169/62/000/010/014/071
D228/D307

onic development is close, can be coordinated with respect to the
values of χ and I_n .

[Abstracter's note: Complete translation]

Card 2/2

VALIYEV, A.A.

Using the residual magnetic method for dividing and correlating
Cenozoic molasses in the Chust-Pap anticline (northern Fergana).
Uzb. geol. zhur. no.2:8-16 '60. (MIRA 13:10)

1. Institut geologii AN UzSSR.
(Fergana--Geology, Stratigraphy)
(Rocks, Sedimentary--Magnetic properties)

VALIYEV, A.A.

Paleomagnetic correlation of the Marguzor section of
Cenozoic continental molasse strata (northern Fergana).

Izv.AN SSSR.Ser.geofiz. no.7:974-976 J1 '60.

(MIRA 13:7)

1. Akademiya nauk Uzbekskoy SSR, Institut geologii i razrabotki
neftnykh i gazovykh mestorozhdeniy.

(Marguzor region--Geology, Stratigraphic)

(Rocks--Magnetic properties)

VALIYEV, A.A.

Determination of the position of the earth's poles during the Tertiary period based on studies of remanent rock magnetism in some regions of northern Fergana. Izv. AN SSSR. Ser. geofiz. no.8:1213-1215 Ag '60. (MIRA 13:8)

1. Institut geologii i razrabotki neftyanykh i gazovykh mestorozhdeniy. (Magnetism, Terrestrial)

24.5300

39129
S/058/62/000/006/069/136
A061/A101

AUTHORS: Bagduyev, G. B., Valiyev, A. A., Kazhlayev, M. A., Kamilov, I. K.

TITLE: The heat conductivity of lead telluride

PERIODICAL: Referativnyy zhurnal, Fizika, no. 6, 1962, 17, abstract 6E143
("Uch. zap. Dagestansk. un-t", 1961, v. 7, no. 1, 107 - 111)

TEXT: The heat conductivity (λ) of PbTe has been measured in the temperature range of 90 - 600°K. A plane stationary method was applied and the measurement accuracy was 4 - 6%. Up to 360°K, $\lambda \sim 1/T$. At higher temperatures the dependence was weaker, which is explained by the influence of the electron contribution to λ . At temperatures higher than 200°K, the measurement results diverge from those of Ye. D. Devyatkova (RZhFiz, 1957, no. 11, 27619), which can be explained by the presence, in the experiments of the latter, of neglected radiation losses from the lateral sample surfaces. ✓

L. Filippov

[Abstracter's note: Complete translation]

Card 1/1

VALIYEV, A.A.; EGAMBERDYEV, M.E., kand. geol.-min. nauk, otv. red.;
TERNOVSKAYA, R.M., red.; KARABAYEVA, Kh.U., tekhn.red.

[Lithology and paleomagnetism of Cenozoic molasses in northern
Fergana] Litologiya i paleomagnetizm kainozoiskikh molass Sever-
noi Fergany. Tashkent, Izd-vo UzSSSR, 1962. 122 p.
(MIRA 15:11)

(Fergana--Rocks, Sedimentary--Magnetic properties)

ACC NR: ARG015870

SOURCE CODE: UR/0275/85/000/012/B007/B008

AUTHOR: Kamilov, I. K.; Kuznetsov, A. A.; Valiyev, A. A.

TITLE: On the investigation of thermoelectric properties of ferromagnetic semiconductors

SOURCE: Ref. zh. Elektronika i yeye primeneniye, Abs. 12B59

REF SOURCE: Sb. aspirantsk. rabot. Dagestansk. un-t. Yestestv. i fiz.-matem. n. Ma: achkala, 1964, 135-146

TOPIC TAGS: ferromagnetic material, nickel base alloy, zinc base alloy, semiconducting material, thermal EMF, resistivity, thermoelectric property

ABSTRACT: Results of measurements of thermal emf and resistance ρ of nickel-zinc ferromagnetic semiconductors of the $Ni_{(x)}ZnO_{(y)}Fe_2O_3(50)$ type of various composition, are given. An investigation is made of specimens the composition of which is determined by the following values of x and y (in mole %): $x = 0-50$, $y = 50-7.5$. In the transition from a zinc ferrite to a nickel ferrite, the values of α increase linearly by a factor of ~ 2 (the measurement temperature is 300K). In the region of 300-600 K, the Kelvin temperature α for all ferrites, except pure nickel ferrites, is negative. The dependence $\alpha(T)$ for the majority of the specimens is weak. In the case when individual ferrites reach the region of magnetic transforma-

Card 1/2

UDC: 539.293:537.32

ACC NR: AR6015870

tion a sharp change in the fluctuation of $\rho(T)$ is observed, which is related to the increasing participation of conductivity electrons in the transitory processes, and also in the change in distances between atoms. The activation energy determined according to $\rho(T)$ increases as ρ increases and is equal to 0.21–0.3 eV for different ferrites. The mobility of the electrons $u = 1/ne\rho$, where the electron concentration n is estimated from α , is equal to 10^{-4} – 10^{-3} cm²/V·sec. The effective mass of carriers is estimated to be equal to 2–7. The methodology of the measurements is described. [Translation of abstract] V. K.

SUB CODE: 09,20

Card 2/2

VALIYEV, A. M.

N. E. Fedoseev - odin iz pervykh marksistov v Rossii [N. E. Fedoseev, one of Russia's first Marxists.] Kazan, Tatgosizdat, 1952. 168 p.

SO: Monthly List of Russian Accessions, Vol. 6, No. 5, August 1953

VALIYEV, A. V. ed.

Sheep breeding 2. izd. ispr. i dop. Moskva, Gos. izd-vo selkhoz. lit-ry, 1952. 287 p.

1. Sheep breeding
2. Sheep - Russia

L 45723-66 ENT(m)/EWP(t)/ETI IJP(c) JD/JG
ACC NR: AP6026418 (A) SOURCE CODE: UR/0249/66/022/001/0012/0015

AUTHOR: Shakhtakhtinskiy, G. B.; Valiyev, B. S.; Aslanov, G. A. 33

ORG: Chemistry Institute (Institut khimii) 61 B

TITLE: Arsenate method of iodometric determination of yttrium in the presence of scandium

SOURCE: AN AzerbSSR. Doklady, v. 22, no. 1, 1966, 12-15

TOPIC TAGS: yttrium, scandium, arsenate, quantitative analysis

ABSTRACT: A new method has been developed for determining yttrium iodometrically as the arsenate in the presence of scandium (at least 0.3 mg of Y in the presence of up to 20 mg of Sc per 25 ml of solution). In order to precipitate the yttrium ion with arsenate ions, heating of the solution should be discontinued immediately after the appearance of turbidity, since further heating leads to the formation of an amorphous precipitate which contains scandium. If such a precipitate does form, 5 to 6 drops of 1 N HCl should be added; this dissolves the amorphous scandium precipitate, but the crystalline yttrium arsenate is not affected and deposits quantitatively. Yttrium as the arsenate is finally determined iodometrically. The analysis lasts about 40 to 50 min. The paper was presented by Academician AN AzerbSSR Nagiyev, M. F. Orig. art. has: 2 tables.

SUB CODE: 07/ SUBM DATE: 09Apr65/ ORIG REF: 005/ OTH REF: 002
Card 1/1 JLR

RASHCHEPKIN, K.Ye.; RAMEYEV, M.K.; VALIYEV, D.M.

Consolidated UIM-14 insulation machine. Transp. i khran. nefti
no.5:8-9 '63. (MIRA 17:3)

1. Nauchno-issledovatel'skiy institut po transportu i khraneniya
nefti i nefteproduktov.

VALIYEV, G.T. (Ufa)

Acquainting students with the work of the "electric shepherd."
Fiz. v shkole 22 no.2:77 Mr-Apr '62. (MIRA 15:11)
(Electric fences)

14(5)

SOV/92-59-2-31/40

AUTHOR: Valiyev, K., Foreman

TITLE: We Will Properly Implement the Resolution Adapted at the Twenty First Congress of the Communist Party of the Soviet Union (Dostoyno vypolnim resheniya XXI s"yezda KPSS)

PERIODICAL: Neftyanik, 1959, Nr 2, pp 31 (USSR)

ABSTRACT: The author states that his drilling crew handles a number of oil wells scattered over a large territory including several administrative districts of the Tatar Republic. Some of these wells are 50-60 km from the oilfield bulk plant and 70-80 km from the town of Al'met'yevsk. This makes the work of the crew extremely difficult. Nevertheless, the discipline, the solidarity, and the high spirit of crew members help to overcome all difficulties, and to gain second place in the All-Union socialist competition contest. The resolution adopted at the Twenty First Congress of the Communist Party encouraged oilmen, stimulated their efforts, and as a result, permitted them to undertake an obligation to produce 2450 tons of petroleum in excess of the target figure set in the production plan. The drilling crew under discussion has also pledged to raise the productivity of labor, to restore four depleted wells, and to maintain the same

Card 1/2

We Will Properly Implement (Cont.)
exploitation ratio.

SOV/92-59-2-31/40

ASSOCIATION: Promysel No 3 NPU Al'met yevneft' (The No. 3 Oilfield of the
Al'met'yevneft' Petroleum Production Administration)

Card 2/2

VALIYEV, K.

Organization and remuneration of machinery operators on collective farms under new conditions. Sots. trud. no.9:64-70 '58. (MIRA 11:10)

(Farm mechanization) (Wages)

Valiyev, K

VALIYEV, K.

Wages of combine operators of machine-tractor stations. Sots. trud
no.2:134-137 F '58. (MIRA 11:1)

1. Zamestitel' nachal'nika Upravleniya normirovaniya truda i zar-
platy Ministerstva sel'skogo khozyaystva RSFSR.
(Machine-tractor stations) (Wages)

VALIYEV, K. A.

56-4-34/54

AUTHOR: Valiyev, K. A.,

TITLE: Magnetic Resonance on Nuclei with Paramagnetic Atoms (Magnitnyy rezonans na yadrakh paramagnitnykh atomov) (Letter to the Editor)

PERIODICAL: Zhurnal Eksperim. i Teoret Fiziki, 1957, Vol. 33, Nr 4, pp. 1045-1047, (USSR)

ABSTRACT: The probability of the nuclear relaxation transitions for the ions Cr^{3+} (odd number of electrons) and the ions V^{3+} (even number of electrons) was calculated. From the calculations follows that in chromium, at the temperature of liquid air, the line width of the nuclear resonance does not exceed some oersted and that the absorption can be observed by modulation of the constant magnetic field. For vanadium the time of the spin lattice relaxation is short and the nuclear resonance can only be observed at low temperatures. Further the probability of the nuclear relaxation transitions in the ethyl sulfates of rare earths was calculated. The mathematical course of calculation is separately published.

ASSOCIATION: Kazan State University (Kazanskiy gosudarstvennyy universitet)

SUBMITTED: June 21, 1957

AVAILABLE: Library of Congress

Card 1/1

VALIYEV, K.A.

Nuclear magnetic resonance in salts of the rare earth elements.
Uch. zap. Kaz. un. 117 no.9:145-148 '57. (MIRA 13:1)

1. Kazanskiy gosudarstvennyy universitet im. V.I. Ul'yanova-Lenina.
Kafedra eksperimental'noy i teoreticheskoy fiziki.
(Rare earth salts) (Nuclear magnetic resonance)

VALIYEV, K.A.

Magnetic resonance spectra and probabilities of magnetic dipole transitions on nuclei of paramagnetic atoms. Uch. zap. Kaz. un. 117 no.9:149-153 '57. (MIRA 13:1)

1. Kazanskiy gosudarstvennyy universitet im. V.I. Ul'yanova-Lenina. Kafedra eksperimental'noy i teoreticheskoy fiziki. (Nuclear magnetic resonance) (Dipole moments)

VALIYEV, K.A., Cand Phys-Math Sci --(diss) "Magnetic resonance
on the nuclei of paramagnetic ions." Kazan', 1958. 6 pp (Min
of Higher Education USSR. Kazan' Order of Labor Red Banner State
Univ in V.I.Ul'yanov-Lenin). 120 copies (KI, 20-58, 92)

AUTHOR: Valiyev, K. A. SOV/126-6-2-1/34
TITLE: Spin-lattice Relaxation and Nuclear Magnetic Resonance
in Salts of Rare Earth Elements (Spin-reshetchnaya
relaksatsiya i yadernyy magnitnyy rezonans v solyakh
redkozemel'nykh elementov)
PERIODICAL: Fizika Metallov i Metallovedeniye, 1958, Vol 6, Nr 2,
pp 193-202 (USSR)

ABSTRACT: The possibility of experimental studies of nuclear
magnetic resonance using the nuclei of paramagnetic
ions of rare earth elements is discussed. It is well
known that the experimental observation of the nuclear
magnetic resonance depends upon the width of the absorption
line. In the magnetically weak crystals of rare earth
salts the line width is largely determined by the spin-
lattice interaction of ions. In the present paper the
author calculates the probabilities of relaxation
transitions between hyperfine sublevels in ions of ethyl
sulphates of rare earths. The calculated probabilities
for relaxation transitions for nuclear spins are compared
with the probabilities for relaxation transitions for
electron spins. For the majority of ions with an odd
number of 4f-electrons the probabilities of nuclear

Card 1/2

SOV/126-6-2-1/34

Spin-lattice Relaxation and Nuclear Magnetic Resonance in Salts
of Rare Earth Elements

relaxation are 100 times smaller than the probabilities in the case of electrons. For ions with an even number of 4f-electrons the probabilities are down by a factor of 10^5 as compared with the electron case. It is shown that in rare earth salts the effect can only be observed at liquid helium temperatures. Explicit formulae are given for the relaxation probabilities of the following ions: Sm^{3+} , Er^{3+} , Dy^{3+} , Tb^{3+} , Ho^{3+} , Tm^{3+} , Yb^{3+} , Eu^{3+} . The work was supervised by S. A. Al'tshuler. There are 2 tables and 11 references, 4 of which are Soviet, 2 German, 5 English.

ASSOCIATION: Kazanskiy gosudarstvennyy universitet
(Kazan' State University)

SUBMITTED: November 24, 1956

Card 2/2

1. Perturbation theory--Mathematical analysis
2. Nuclear spins
3. Rare earth salts--Temperature factors
4. Ethylsulfates--Properties

SOV/126-6-5-2/43

AUTHOR: Valiyev, K. A.

TITLE: On Magnetic Resonance of Nuclei of the Iron-Group Atoms in Ionic Crystals (O magnitnom rezonanse v ionnykh kristallakh na yadrakh atomov elementov gruppy zheleza)

PERIODICAL: Fizika Metallov i Metallovedeniye, 1958, Vol 6, Nr 5, pp 776-780 (USSR)

ABSTRACT: The author discusses nuclear and electron magnetic resonances of paramagnetic vanadium and chromium atoms in ionic salts. It was found that even at temperatures of the order of 1°K the electron spin relaxation contribution (A_e) predominates over the nuclear relaxation part (A_N) in the expression for the total width of a magnetic resonance line: $\gamma = A_e + A_N$. Calculations showed that, in chromium salts at 2°K, A_e/A_N for Cr^{3+} ions is of the order of 10^4 . For V^{3+} ions in vanadium salts the same ratio is of the order of 10^5 at 2°K. A table on p 780 gives the calculated values of the electron (τ_e , in μ sec) and nuclear (τ_N , in sec) relaxation time constant at 90, 64.4 and 2.04°K of Cr^{3+} ions in chrome alums. These values indicate that

Card1/2

On Magnetic Resonance of Nuclei of the Iron-Group Atoms in Ionic Crystals

SOV/126-6-5-2/43

nuclear magnetic resonance in chromium (spin $S=1$) and vanadium ($S=3/2$) may be observed only at liquid-helium temperatures. The paper is entirely theoretical. Acknowledgment is made to S. A. Al'tshuller who directed this work. There are 2 figures, 1 table and 8 references, 2 of which are Soviet, 4 English, 1 Dutch and 1 translation.

ASSOCIATION: Kazanskiy gosudarstvennyy universitet
(Kazan' State University)

SUBMITTED: January 14, 1957

Card 2/2

AUTHORS: Valiyev, K. A., Bashkirov, Sh. Sh. SOV/56-35-1-54/59

TITLE: A Stimulated Amplifier of Radio-Frequency Signals on the Basis of the Hyperfine Sublevels of Paramagnetic Atoms (Stimulirovanny usilitel' radiochastotnykh signalov na sverkh-tonkikh podurovnyakh paramagnitnykh atomov)

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1958, Vol. 35, Nr 1, pp. 302 - 303 (USSR)

ABSTRACT: The idea of applying paramagnetic crystals as working piece of stimulated amplifiers for centimeter waves was conceived by N. Bloembergen (Blombergen) (Ref 1). Such an amplifier was constructed on the basis of the salts of the Gd^{3+} ion, using the dipole-transitions between the sublevels of the energy of the electron spins of the paramagnetic Gd^{3+} ions. Some previous papers concerning this subject are mentioned. The authors of this paper wish to draw attention to the possible amplification of 10^8 - 10^9 Hertz signals by application of the transitions between the hyperfine sublevels of the paramagnetic ions. As an example, the crystals of the salts of the divalent ions of Cu^{64} (ground state 2D , $S = 1/2$, $I = 1$) are investigated. The con-

Card 1/2

A Stimulated Amplifier of Radio-Frequency Signals SOV/56-35-1-54/59
on the Basis of the Hyperfine Sublevels of Paramagnetic Atoms

clusions obtained in this paper seem to be applicable also to other paramagnetic atoms. The scheme of the spin sublevels of Cu^{++} in a strong magnetic field is shown in a figure, and also the numerical values of the probabilities of the relaxation transitions are given. In pulsed schemes with pulses of 10^{-4} sec, the power yield may attain values of $\sim 10^{-3}$ W. The authors thank S.A.Al'tshuler for the discussion of results. There are 1 figure and 6 references, 1 of which is Soviet.

ASSOCIATION: Kazanskiy pedagogicheskiy institut (Kazan' Pedagogical Institute)

SUBMITTED: April 15, 1958

Card 2/2

24(5)

SOV/56-35-3-18/61

AUTHORS:

Bashkirov, Sh. Sh., Valiyev, K. A.

TITLE:

On the Polarization of the Nuclear Moments and the Width of Nuclear Resonance Lines in Crystals of Cu^{++} -Ion Salts (0 polyarizatsii yadernykh momentov i shirine liniy yadernogo rezonansa v kristallakh soley iona Cu^{++})

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1958, Vol 35, Nr 3, pp 678-684 (USSR)

ABSTRACT:

In the introduction several papers dealing with this subject, which have already been published (Refs 1-7), are discussed (as e.g. on the investigation of the degree of polarization of the magnetic moments at 1 - 4°K, experiments carried out concerning the polarization of nuclei with the aid of electron transitions; Feher (Fekher) investigated the nuclear polarization of paramagnetic atoms; investigation of transition probabilities (Refs 5-7)). The present paper aims at calculating the probabilities of nuclear relaxation transitions in Cu^{++} -ion salts $a_{m,m-1}$ and the probability of electronic transitions $A_{M,M-1}$ (M = magnetic quantum number of electron spin) (such

Card 1/3

SOV/56-35-3-18/61

On the Polarization of the Nuclear Moments and the Width of Nuclear Resonance Lines in Crystals of Cu^{++} -Ion Salts

calculations of nuclear- and electronic relaxation transitions for $S = 1/2$ and $I = 1/2$ have already been carried out by Abraham (Abraham) (Ref 2)). The amount of nuclear polarization occurring as a result of the saturation of electron- and nuclear resonance transitions is calculated. The chapters of this paper deal with the following subjects: 1) Introduction; 2) Probabilities of nuclear relaxation transitions (ansatz for transitions between hyperfine sublevels: $\mathcal{K} = \mathcal{K}_{OL} + \lambda(\vec{L}\vec{S}) + A I_z S_z + B(I_x S_x + I_y S_y)$; $\lambda(\vec{L}\vec{S})$ - spin-orbital interaction, \mathcal{K}_{OL} - an operator, ansatz for H_{OL} according to Kronig (Ref 8)); 3) Polarization of Cu^{64} nuclei in paramagnetic salts; ($I = 1$) 4) On the breadth of paramagnetic resonance lines on the nuclei of paramagnetic atoms - the effect of electron and nuclear relaxations on the broadening of lines is considered. There are 1 figure and 15 references, 5 of which are Soviet.

Card 2/3

SOV/56-35-3-18/61
On the Polarization of the Nuclear Moments and the Width of Nuclear Resonance
Lines in Crystals of Cu^{++} -Ion Salts

ASSOCIATION: Kazanskiy gosudarstvennyy pedagogicheskiy institut (Kazan'
State Pedagogical Institute)

SUBMITTED: March 2, 1958

Card 3/3

24(3)

AUTHORS:

Al'tshuler, S. A., Valiyev, K. A.

SOV/50-35-4-1/19

TITLE:

On the Theory of Longitudinal Relaxation in Liquid
Solutions of Paramagnetic Salts (K teorii prodol'noy
relaksatsii v zhidkikh rastvorakh paramagnitnykh soley)

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1958,
Vol 35, Nr 4, pp 947 - 958 (USSR)

ABSTRACT:

In their introduction the authors discuss the papers
already published and dealing with the theory of
longitudinal (spin-lattice) relaxation in para-
magnetic salt solutions; in the USSR B.M.Kozyrev
(Ref 4) is occupied with this problem. The present
paper deals with theoretical investigations in salt
solutions of elements of the groups of iron and rare
earths. Calculations are based on the assumption
that, like in the case of ion crystals, the liquid
solution contains paramagnetic complexes. The normal
vibrations of these complexes are disturbed by the
Brown (Broun) motion; this causes variation of the

Card 1/3

On the Theory of Longitudinal Relaxation in Liquid
Solutions of Paramagnetic Salts

SOV/56-35-4-17/52

electric field in which the paramagnetic ions occur. This variation, in turn, influences the spin-orbital interaction of the electrons of the paramagnetic ion, and may lead to a re-orientation of its magnetic moment, causing relaxation transitions between various steady states. It is shown that, if the interval between the low Stark (Shtark) ion sublevels is $\delta \approx 2kT$, there exist two relaxation times, which are caused by 1) transitions between various Stark (orbit)-levels without changing the direction of spin, and 2) transitions within a Stark level with changed spin orientation. The breadth of the paramagnetic resonance line is due to transitions of the first type. Theoretical results are in good agreement with experimental data concerning copper-, chromium-, and manganese salt solutions. In conclusion, the authors thank B.M.Kozyrev for discussing results. There are 3 figures and 21 references, 11 of which are Soviet.

Card 2/3

On the Theory of Longitudinal Relaxation in Liquid
Solutions of Paramagnetic Salts

SOV/56-35-4-17/52

ASSOCIATION: Kazanskiy gosudarstvennyy universitet (Kazan' State
University) Kazanskiy pedagogicheskiy institut (Kazan'
Pedagogical Institute)

SUBMITTED: March 18, 1958

Card 3/3

SOV/56-36-6-18/66

24(2)

AUTHOR: Valiyev, K. A.

TITLE: Contribution to the Theory of Lattice - Spin Relaxation (K teorii reshetchnoy spinovoy relaksatsii)

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1959, Vol 36, Nr 6, pp 1743 - 1749 (USSR)

ABSTRACT: This article gives a theoretical analysis of the relaxation of systems containing two kinds of spins which influence each other and have widely differing relaxation periods. For this purpose the method of correlation functions for rapidly variable spin values of the system is used. In the introduction to the present article reference is made to the following papers: A paper by J. Waller on the theory of relaxation in the magnetic interaction of particles of cubic crystals, a paper by N. Bloembergen on the relaxation of dipole-like interacting particles, which is of particular importance for the present paper because it is shown that Bloembergen's theory also describes the processes of spin relaxation in molecules of diamagnetic crystals, a paper by S. A. Al'tshuler on

Card 1/3

Contribution
to the Theory of Lattice - Spin Relaxation

SOV/56-36-6-18/66

lattice relaxation in liquids, especially of compounds of the form XY_6 , where X denotes a paramagnetic ion, Y - a molecule of the solvent or an anion. The present paper describes the relaxation process for a kind of spin which is connected with exchange- or dipole forces with spins of a different kind. Determination of the latter is, however, carried out independently of the former, as in their case the relaxation periods are very small. Experiments concerning the hyperfine structure of the spectra of magnetic nuclear resonance confirm the existence of an interaction between the quadrupole moment of the nuclei with the electric field of the surrounding nuclei. An especially interesting example is that of the relaxation of the nuclear spins of paramagnetic atoms caused by hyperfine interactions with the spins of the electron shell of the same atom. There follows an analysis of rapidly variable spin systems, and reference is made to a paper by A. I. Rivkind, which explains the relaxation of protons in liquid solutions of paramagnetic salts. In this connection, the variation of the quantization of electron spins with respect to the external statistical magnetic field plays an important part. Calcula-

Card 2/3

Contribution
to the Theory of Lattice - Spin Relaxation

SOV/56-36-6-18/66

tions concerning the quadrupole broadening of the lines of proton resonance in liquids follow. Also resonance phenomena in spin-lattice relaxation as well as the lattice relaxation of nuclei belonging to paramagnetic atoms are analyzed. The author finally thanks Professor S. A. Al'tshuler and Professor B. M. Kozyrev for discussions. There are 17 references, 4 of which are Soviet.

ASSOCIATION: Kazanskiy pedagogicheskiy institut (Kazan' Pedagogical Institute)

SUBMITTED: December 8, 1958

Card 3/3

SOV/56-37-1-17/64

21(1)

AUTHOR:

Valiyev, K. A.

TITLE:

The Theory of Spin-lattice-quadrupole Relaxation of Nuclei in Liquid Solutions of Diamagnetic Salts (Teoriya spin-reshetochnoy kvadrupol'noy relaksatsii yader v zhidkikh rastvorakh diamagnitnykh soley)

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1959, Vol 37, Nr 1(7), pp 109-117 (USSR)

ABSTRACT:

The calculations were carried out under the assumption that stable complexes originate in the solutions around the ions. These complexes consist of the molecules of the solvent, or simultaneously of molecules of the solvent and of anions (or cations). The first part deals with the mechanism of spin-lattice relaxation in diamagnetic ion solutions. The ion to be investigated is assumed to be in the center of a mixed complex, which e.g. partly consists of water molecules and partly of anions. In the center of the mixed complex, a strong inhomogeneous electric field is assumed to exist there which causes a quadrupole separation of the spin levels of the ion nucleus. This separation is superimposed over the Zeeman separation. Under real conditions, mixed and not

Card 1/3

SOV/56-37-1-17/64

The Theory of Spin-lattice-quadrupole Relaxation of Nuclei in Liquid Solutions of Diamagnetic Salts

mixed complexes are present in the solution at the same time. In the next parts, the following factors are calculated and discussed: the energy of the interaction of the lattice and of the spins of the nuclei with an electric dipole moment, the probability of relaxation transitions, the relaxation time T_1 , and the width of lines, the quadrupole relaxation of the nuclear spin caused by a diffuse rotation of a mixed complex. The author arrives at the following conclusions:

1) The time of spin-lattice relaxation of the spins of the nuclei with large electric quadrupole moment in mixed complexes is in the order of magnitude of 10^{-6} to 10^{-7} sec. This corresponds to a line width of $\sim 10^6$ cycles. A line of such a width evidently cannot be observed by experiment. Therefore, the intensity of experimentally observable lines of the magnetic resonance on nuclei with a high quadrupole moment is proportional to the number of complexes only consisting of water. 2) For aqueous complexes of the ions Al^{3+} and Ga^{3+} , a good agreement is obtained between the measured and calculated values of the resonance line widths. It is known from chemical investigations that the hydrate complexes of the

Card 2/3

SOV/56-37-1-17/64

The Theory of Spin-lattice-quadrupole Relaxation of Nuclei in Liquid Solutions of Diamagnetic Salts

ions Al^{3+} and Ga^{3+} are very stable. This confirms the correctness of the model used for calculations in the present paper if only the life-time of the complexes is longer than the spin-lattice relaxation of the nuclei. 3) The calculated and the experimentally observed value of the resonance line width do not agree in the case of the ions Na^+ , Br^- , J^- . This is apparently connected with the weak stability of the complexes formed by the ions Na^+ , Br^- and J^- with water: The experimentally observed line width of the nuclei Na^{23} , Br^{79} , Br^{81} and J^{127} is not connected with the time of spin-lattice relaxation but with the life-time of the complex itself. There are 1 table and 15 references, 6 of which are Soviet.

ASSOCIATION: Kazanskiy gosudarstvennyy pedagogicheskiy institut
(Kazan' State Pedagogical Institute)

SUBMITTED: January 10, 1959

Card 3/3

83732

S/056/60/038/004/025/048
B006/B056

24.6520
AUTHOR:

Valiyev, K. A.

TITLE:

The Theory of the Quadrupole Relaxation of Nuclear Spins in Liquids

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1960,
Vol. 38, No. 4, pp. 1222 - 1232

TEXT: The present paper theoretically investigates the line width of the nuclear-magnetic resonance in liquids, in which the atoms investigated have a quadrupole moment and form no stable complexes with other particles of the liquid. These conditions are e.g. realized in the investigation of the quadrupole resonance of nuclear spins of halide ions in aqueous electrolyte solutions. The water molecules may (with known accuracy) be considered to be point charges. The calculated line width is influenced by the selection of the effective molecule charge, but not the physically essential dependence of the line width on the temperature and nature of the thermal particle motion. Analogous conditions prevail also in the melts of metals and ion salts. The author assumes the thermal

Card 1/3

83732

The Theory of the Quadrupole Relaxation of
Nuclear Spins in Liquids

S/056/60/038/004/025/048
B006/B056

motion of the liquid particles to be a free translational diffusion. This assumption is satisfied best for liquids at a temperature that is considerably higher than the melting point. It is in such a liquid that a single ion is now investigated, the nucleus of which has an electric quadrupole moment. The remaining particles are considered to be point charges which are chaotically distributed around the ion investigated, so that an inhomogeneous resulting E-field acts upon the nucleus of this ion. The relative position of the individual particles changes continuously due to diffusion, so that the E-field acting upon the one ion is a random time function. The E-field influences its spin via the quadrupole moment of the nucleus; the consequence is a random shifting of the sublevels of nuclear spin, so that a secular broadening of resonance lines and a broadening caused by relaxation transitions between the sublevels occurs. First, the theoretical method is explained and expressions are obtained for the energy of the quadrupole moment of the nucleus in the field of the point charges. The next parts of the paper deal with the correlation function and the relaxation times T_2 and T_1 (T_2 characterizes the secular line broadening, T_1 - the non-secular line broadening caused by a

Card 2/3

83732

The Theory of the Quadrupole Relaxation of
Nuclear Spins in Liquids

S/056/60/038/004/025/048
B006/B056-

shortening of the lifetime of the spin in a certain state because of relaxation transitions between the states. T_2 and T_1 are defined by (6) and (11). For the two times formulas (28) and (32)¹ are then obtained, after which the special case of aqueous electrolyte solutions is studied and numerical calculations and comparisons are discussed. T_2^{-1}/T is obtained (- viscosity of the liquid); this is in good agreement with measurements of T_2^{-1} for the I^{127} nuclear spin in aqueous solutions of NaI and KI salts. Also the conditions in metal melts and melts of ion salts are discussed, and numerical measuring results are compared with theoretical considerations. S. A. Al'tshuler, G. V. Skotskiy, and A. A. Kokin are mentioned. There are 18 references: 6 Soviet, 7 US, 1 Japanese, and 4 British.

ASSOCIATION: Fiziko-tekhnicheskiy institut Kazanskogo filiala Akademii nauk SSSR (Institute of Physics and Technology of the Kazan' Branch of the Academy of Sciences, USSR)

SUBMITTED: October 23, 1959
Card 3/3

VALIYEV, K.A.

Theory of the line widths of vibration and Raman spectra of
molecules in dipole liquids. Opt. i spektr. 11 no. 4: 465-470
0 '61. (MIRA 14:10)

(Molecular spectra)

VALIYEV, K.A.; KHABIBULLIN, B.M. (Kazan')

Nuclear magnetic resonance and structure of aqueous solutions of electrolytes. Zhur.fiz.khim. 35 no.10:2265-2274 0 '61. (MIRA 14:11)

1. Kazanskiy pedagogicheskiy institut.
(Electrolyte solutions)

VALIYEV, K.A.

Theory of the dissipation of the energy of molecular oscillations in fluids. Zhur. eksp. i teor. fiz. 40 no.6:1832-1837 Je '61. (MIRA 14:8)

1. Kazanskiy pedagogicheskiy institut.
(Molecular dynamics)

28756

S/056/61/041/003/009/020
B125/B102

24.2110 (1138, 1147, 1164)

AUTHORS: Valiyev, K. A., Zaripov, M. M.

TITLE: Theory of spin-lattice relaxation in liquid solutions of electrolytes

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 41, no. 3(9), 1961, 756-761

TEXT: The spin-lattice relaxation of hydrated metal ions in aqueous solutions, caused by interaction of the ionic spin with the internal oscillations of the complex, has been studied by the methods of the theory of random processes. The energy of interaction of the spin \vec{S} of the central ion of the complex with the oscillations Q_j can usually be represented as a series: $\mathcal{H} = \mathcal{H}^{(1)} + \mathcal{H}^{(2)} + \dots = \sum_j P_j(\vec{S}) Q_j + \sum_{ij} P_{ij}(\vec{S}) Q_i Q_j + \dots$ (1)

where $P_j(\vec{S})$ and $P_{ij}(\vec{S})$ are quadratic functions of the components of the vector $\vec{S}(S_x, S_y, S_z)$. S. A. Al'tshuler and K. A. Valiyev (ZhETF, 35, 947, 1958) calculated the relaxation rate of the spin \vec{S} , which is due to the

Card 1/7